Delaware Public Health Laboratory State CRE Laboratory

CDC Epidemiology & Laboratory Capacity (ELC)
Grant

Debbie Rutledge, MBA MT(ASCP) Infectious Disease Laboratory Manager II

Antimicrobial Resistance

- 2 million illnesses & 23,000 deaths annually related to Antibiotic Resistance (AR)
- Carbapenem-Resistant Enterobacteriaciae (CRE) increasing resistance to all theraputic agents.
- Increasing resistance of *Pseudomonas aeruginosa* (intrinsic, non carbapenemase vs. carbapenemase mechanisms)
- Carbapenemases = transferable = greater public health threat

State CRE Laboratory

- Develop capacity to confirm CRE & carbapenem resistant *Pseudomonas* aeruginosa (CRPA) & detect carbapenemase mechanisms.
- Early detection & implement timely local prevention efforts
- Improve response to outbreaks & speed up identification of CRE
- Antimicrobial Resistance Laboratory Network with 7 regional labs for enhanced bacteria and fungal testing capabilities (Maryland).

ELC Grant

- New activity just awarded August 30, 2016
- ▶ First conference call with CDC 9/30/2016
- DPHL to participate in HAI/AR Prevention programs with hospital infection control preventionists (ICP)
- Adopt & validate PCR and other testing methods to detect most common & important resistance mechanisms
- Implement bacterial isolate storage, transport to AR Regional labs as requested.


Recommended Laboratory testing

- Species Identification MADITOF or automated instrument
- Antimicrobial Susceptibility testing by another method than provider-disc diffusion/Etest or broth microdilution
- Phenotypic methods for carbapenemase production-carbapenem inactivation method or CarbaNP assay
- Molecular detection PCR targets
 - bla KPC & bla NDM-1 genes
 - OXA-48 like genes
 - bla VIM
 - Other optional targets bla IMP, & mcr 1

Collect Isolates from HCF

- All Escherichia coli, Klebsiella oxytoca, Klebsiella pneumoniae, & Enterobacter spp.
 - Resisitant to imipenem, meropenom, doripenem
 (MIC ≥ 4ug/ml) or ertapenem (≥ 2)
- All Pseudomonas aeruginosa
 - Resistant to imipenem, meropenem, or doripenem
 (≥ 8)
- If states do not have the capacity to test all CRE isolates, may choose sampling scheme based on burden & local testing capacity

CDC Antibiotic Resistance Laboratory Network: 7 Regional Labs



Questions or Comments

- Workload Volume? Can hospital laboratories estimate # isolates from last year?
- We need your help in sending us these targeted isolates.
- More information can be found at
- http://www.cdc.gov/drugresistance/solutions -initiative/ar-lab-networks.html